

HPAI A(H5) Scenario-Based Human Health Risk Assessment



Summary: HPAI A(H5) Scenario-Based Human Health Risk Assessment for the United States as of October 3, 2024

Based on the latest information from the Missouri human H5N1 case, the Center for Outbreak Response Innovation (CORI) judges that the ongoing H5N1 outbreak in cattle, poultry, and other animals is still within Scenario 3.

On September 6 CDC reported the [first human case of H5N1 bird flu in Missouri](#), the first and only reported human case in the ongoing outbreak with no known animal exposure and symptomatic contacts. On September 20, CDC reported that there was a [third symptomatic contact](#) connected to the Missouri case. This individual is a healthcare worker that cared for the patient while they were hospitalized. The third contact did not receive testing for influenza. On September 27, CDC reported that a total of [18 healthcare workers had high risk exposure and 94 healthcare workers had low risk exposure](#) to the case while they were hospitalized. From these exposures, four additional symptomatic contacts have been identified. Although all ill individuals have recovered, there remains concern that the source of the infection for the case patient is still unknown and that human-to-human transmission from the original case to the healthcare worker(s) is possible and may have been undetected. CDC reports that the original case, the household contact, and the symptomatic healthcare worker contacts have had specimens collected for serologic testing to look for antibodies to avian influenza A(H5) virus, which would indicate a previous infection. No further information has been shared regarding other potential source of H5N1 transmission for the case patient. There is still significant concern that the recovered case is a member of the general public, which could indicate there are chains of transmission that are not currently identified, although the case is currently described by CDC as a isolated event. There is also concern about the potential for human-to-human transmission if the symptomatic healthcare workers are positive for H(5) antibodies, although CDC maintains the likelihood of efficient transmission among humans remains low based on the most recent viral genomic sequence. We are not raising the risk level at this time because these additional chains of transmission are not confirmed and transmission from the case to the healthcare workers has not been confirmed, however, due to the limited information our confidence in the current risk level is low. *See a detailed analysis beginning on the next page.*

	Risk to farm workers	Risk to other people in contact with affected workers and animal populations	Risk to healthcare workers	Risk to the US general public
Scenario 3 – Increased potential for reassortment and human adaptation, still no human-to-human transmission	Moderate-High	Moderate	Low	Low

Our **confidence** in these risk scores remains low, due to limited information regarding the transmission route for the Missouri case, the uncertainty of potential transmission to healthcare workers, and minimal testing in cattle, humans, and other animals. We recommend 1) continued and enhanced surveillance and transparency regarding potential transmission routes and serology for the Missouri case investigation, 2) continued and enhanced [surveillance for avian influenza](#), and 3) [continued and enhanced precautions for people in contact with birds and other farm animals](#).

HPAI A(H5) Scenario-Based Human Health Risk Assessment



HPAI A(H5) Scenario-Based Human Health Risk Assessment for the United States Center for Outbreak Response Innovation (CORI) Updated as of October 3, 2024

Since the last update on September 17, 2024:

- [On September 20](#), CDC reported that a second healthcare worker that cared for the patient while they were hospitalized had developed symptoms and not been tested.
- [On September 27](#), CDC reported that the two symptomatic healthcare workers reported previously had been exposed prior to following droplet precautions (ie high risk exposure) and that an additional 16 healthcare workers exposed to the case before droplet precautions were instituted. CDC reported that an additional 94 healthcare workers were also exposed to the case after droplet precautions were followed (ie low risk exposure).
- [Four additional ill healthcare worker contacts were reported by CDC](#) on September 27, bringing the current total to seven symptomatic contacts. One of the additional contacts had a high risk exposure while three of the additional contacts had a low risk exposure.
- Of the [seven total symptomatic contacts](#), only one had tested for flu and was subsequently negative. CDC reports that the original case, the household contact, and the symptomatic healthcare worker contacts have had specimens collected for serologic testing to look for antibodies to avian influenza A(H5) virus, which would indicate a previous infection.
- CDC and CORI maintain that the risk to the general public remains low at this time. The risk to farmworkers remains moderate-high and the risk to other people in contact with affected workers and animal populations remains moderate. The confidence in these risk scores is low due to limited information on a potentially new chain of transmission, the limited information on the health status for the healthcare workers and the limited testing in humans, dairy cows, and other animals. CORI may adjust this assessment based on new information about the chain of transmission for the Missouri case, new results of ongoing serology testing, or identification of additional human cases with no known animal exposures through routine influenza surveillance.

Highly pathogenic avian influenza (HPAI) A(H5N1) clade 2.3.4.4b viruses began circulating in Europe, Africa, and Asia in 2020, leading to an unprecedented number of deaths in wild birds and poultry. In 2021 and 2022, the virus spread across the globe, reaching North, South, and Central America. In 2023, the Food and Agriculture Organization of the United Nations (FAO), the World Health Organization (WHO), and the World Organization for Animal Health (WOAH) released a [joint report](#) regarding the increasing cases of land and sea mammals infected with A(H5N1) clade 2.3.4.4b, including outbreaks with high mortality. The [US Department of Agriculture Animal and Plant Health Inspection Service](#) (USDA APHIS) began tracking the ongoing panzootic in wild birds and poultry in the US in 2022, and now reports detections [in nearly 10,000 wild birds and over 100 million poultry](#). In March 2024, A(H5N1) clade 2.3.4.4b was detected for the first time in [dairy cattle](#)

HPAI A(H5) Scenario-Based Human Health Risk Assessment



in the US. Since then, the virus has been detected in nearly 200 cattle herds across more than one-fifth of US states and more than a dozen farm workers have tested positive for H5N1. The majority of H5 [human cases](#) associated with H5N1 outbreak in the US have been among farm workers with direct exposure to poultry and cattle ([four from affected cows and nine from affected poultry](#)), although one human H5 case has [no known animal exposure](#). The states have reported cases include [Texas](#), [Michigan](#), and [Colorado](#), and [Missouri](#).

To date, 13 of the 14 human cases in the current US outbreak have been mild, with conjunctivitis (eye redness/pink eye) and mild respiratory symptoms being most common. One case with underlying conditions had chest pain, nausea, vomiting, diarrhea, and weakness. None of the cases have resulted in severe illness managed in the ICU. All cases have been treated with antivirals and have recovered. There has been no documented human-to-human transmission at this time. Based on sequenced human samples, CDC currently reports there are [no markers of antiviral resistance](#). Sequencing largely indicated that the candidate vaccine viruses (CVVs) are [expected to provide protection](#); however, amino acid changes in the one Missouri case may impact cross reactivity of CVVs, although additional antigenic testing is ongoing.

Notably, however, the risks to human health from this outbreak are highly uncertain because of a current lack of surveillance data and other basic scientific and epidemiological information. The situation is complex and may change rapidly. Risk assessment can be very helpful in times of significant uncertainty because it enables structured consideration of complex scenarios, likelihoods, and consequences to inform decisions around policy and operational action, as well as implementation of protective measures and future planning for worst-case scenarios. It is important not to wait for perfect information to estimate potential risk, because decisions must be made even in the absence of plentiful data.

Therefore, the Center for Outbreak Response Innovation (CORI) conducted a scenario-based risk assessment to consider human health risks both now and in potential future scenarios. We will update this assessment as additional data become available.

***Please note:** We are evaluating the risks to human health should each scenario occur, **not** the relative risk of any one scenario occurring.

Features that would characterize each scenario include:

Scenario 1 – Minimal spread in cattle: The virus is predominantly infecting cattle and there is minimal spread within herds and to other animals. Likelihood of widespread human infections is low. Population health consequences are low. Overall risk to human health in this scenario is low.

Scenario 2 – Widespread transmission in cattle, few human infections, no human-to-human transmission: The virus is predominantly infecting cattle but spreads widely within herds. There is also occasional cow-to-human transmission but no human-to-human

HPAI A(H5) Scenario-Based Human Health Risk Assessment



transmission. Likelihood of widespread human infections is low. Population health consequences are low. Overall risk is low, but population-specific risk is increased for farm workers.

Scenario 3 – Increased potential for reassortment and human adaptation, still no, or very limited, human-to-human transmission: The virus begins to infect swine or other animal species that could facilitate the mixing and spreading of influenza viruses. This increases the likelihood that the virus reassorts with other influenza viruses and adapts to humans. In this scenario, we expect that some limited human-to-human transmission could be reported but only among farm workers or close contacts of those workers, and not among healthcare workers. Likelihood of widespread human infections is low. Population health consequences are low. Overall risk of widespread transmission in humans is low, but risk is increased for farm workers and close contacts of those workers. The relative risk of a future pandemic has increased, but the absolute risk remains low.

Scenario 4 – Increasing reports of human infections, limited human-to-human transmission between close contacts: There are more reports of human infections due to contact with infected animals like cattle, swine, and/or poultry. Limited human-to-human transmission is reported among close contacts of infected individuals, including healthcare workers, but there is no efficient human-to-human transmission. Likelihood of widespread human infections is moderate. Population health consequences are low. Overall risk of widespread transmission is low, but population-specific risk is increased for farm workers, close contacts of farm workers, and healthcare workers. The likelihood of a future pandemic is increased.

Scenario 5 – Efficient human-to-human transmission: There are reports of efficient human-to-human transmission. Likelihood of human infections is high because the virus now transmits efficiently and will be very difficult to contain. Population health consequences are high. Overall risk is high for all populations. The likelihood of a pandemic is very high.

Due to recent events, CORI judges that the ongoing H5N1 outbreak in cattle and poultry is now within Scenario 3, meaning the virus is infecting multiple animal species that facilitate the mixing and spreading of influenza viruses, increasing the likelihood that the virus reassorts with other influenza viruses and adapts to humans.

This judgment is based on the widespread occurrence of H5N1 infections in cattle herds across more than one-fifth of US states since March 2024, continued reports of human H5 among dairy workers and poultry workers, continued reports from USDA APHIS of H5N1 in other species (domestic cats, racoons, foxes, mice, and others), and new information from [USDA](#) and early access articles in *Nature* regarding the potential for [cow-to-cow and cross-species transmission](#) and the potential for H5N1 to bind with sialic acids expressed in [human respiratory upper airways](#). This assessment also builds on the previous judgment regarding the detection of H5 in wastewater

HPAI A(H5) Scenario-Based Human Health Risk Assessment



in [multiple states](#), high mortality among H5N1-infected cats that live on [affected farms](#), and remaining concern from a preprint article that cows may serve as mixing vessels for influenza reassortment because of the existence of both avian and human influenza [receptors in cows](#).

Although all studies regarding the potential for cows as mixing vessels and cow-to-cow and cross-species transmission are preprint and early access, respectively, meaning they have not been either peer-reviewed, finalized, or replicated, the continued transmission of H5N1 from poultry and cattle to humans in combination with the recent preliminary findings from epidemiologic studies raise concern for increased reassortment potential. To date, no human-to-human transmission has been reported in the current outbreak, however, recent events may increase the potential for transmission to close contacts of affected farm workers and animal populations.

At this time, CORI judges that the likelihood of widespread human infections is low and the population health consequences are low. Furthermore, the overall risk of widespread transmission in humans remains low, but risk is increased for farm workers and close contacts of farm workers. The relative risk of a future pandemic has increased compared to the previous months, but the absolute risk remains low.

CORI are continuing to monitor and adapt this risk assessment to the evolving changes of the H5N1 outbreak in the US. Factors that may contribute to a change in this assessment include increased reports of H5N1 human infections among farm workers, reports of human-to-human transmission, evidence of H5N1 clade 2.3.4.4b adapting for human-to-human transmission, or detection of H5N1 among individuals in the US without a known exposure to infected animals.

H5N1 Human Health Risk Assessment Scenario Table

	Risk to farm workers	Risk to other people in contact with affected workers and animal populations	Risk to healthcare workers	Risk to the US general public
Scenario 1 – Minimal spread in cattle	Low	Low	Low	Low
Scenario 2 – Widespread transmission in cattle, few human infections, no human-to-human transmission	Moderate	Low	Low	Low
Scenario 3 – Increased potential for reassortment and human adaptation, still no, or very limited,	Moderate-High	Moderate	Low	Low

HPAI A(H5) Scenario-Based Human Health Risk Assessment



human-to-human transmission				
Scenario 4 – Increasing reports of human infections, limited human-to-human transmission between close contacts	High	Moderate-High	Moderate	Low-Moderate
Scenario 5 – Efficient human-to-human transmission	High	High	High	High

Methods: The purpose of this document is to consider possible future developments in this outbreak and describe corresponding risks to human populations should a given scenario occur. A risk score is determined for each key population across multiple possible scenarios. Risk scores are evaluated on a five-point scale, with scores including low, low-moderate, moderate, moderate-high, and high. The confidence in each of these risk level assignments are based on the breadth, depth, and the quality of information available. The overall confidence of each risk score determination is based on a three-point scale, which includes low, moderate, and high. Each risk score and the confidence in the score is discussed in the Appendix.

In each scenario regarding H5N1, we consider the risk to four distinct populations: farm workers on affected farms, other people in the vicinity of affected animal populations and farm workers (eg, household contacts of workers, people living near affected facilities with potential contact with infected animals or farm workers, healthcare providers treating infected individuals), US healthcare workers, and the US general public.

In determining these risks, we consider several factors, including cow-to-cow transmission pathways (eg, respiratory transmission, drinking contaminated water and feeding on contaminated grasslands, aerosolization of the virus through the milking process, etc.), cross-species transmission (eg, cow to domestic cat, bird, racoon, etc.), cow-to-human transmission pathways (eg, unprotected and close contact with infected animals, consumption of unpasteurized dairy products, etc.), and human-to-human transmission pathways (eg, aerosol, oral, direct contact). We also consider disease morbidity and mortality, instances of transmission, the level of testing conducted in cattle and other farm animals, the level of testing conducted in humans, existing processes to limit spread from infected animals, genomic surveillance capabilities, and any new mutations showing greater potential for sustained human-to-human transmission. Other factors include events that could increase human-to-human transmission (eg, mass gatherings, seasonal trends, school terms, etc.), treatments available to humans (eg, antivirals), preventative measures against animal-to-worker transmission (eg, use of N95 mask or equivalent, goggles, gloves, gown, head cover, and boot covers), preventative medical countermeasures (eg, vaccines), preventative

HPAI A(H5) Scenario-Based Human Health Risk Assessment



nonpharmaceutical interventions for human-to-human transmission (eg, isolation of infected people, mask wearing), and ongoing response operations to address the outbreak.

Appendix

Scenario 1: H5N1 outbreak stays predominantly in cattle and has minimal spread within herds and to other animals.

In the first scenario, we considered the risk to human health if the H5N1 virus stays in cattle and has minimal spread within herds and to other animals. We determined the health risk to **farm workers** to be **low**, the health risk to **other people in the vicinity of affected workers and animal populations** to be **low**, the risk to **healthcare workers** to be **low**, and the health risk to the **US general public** (and the consequent risk of a pandemic) to be **low**.

Our **confidence** in these risk scores is high given the current level of information known for each of these factors and the fact that the virus is not actively spreading to humans or within human populations in this scenario. To minimize the spread of H5N1 in animals, [USDA](#) recommends:

- Increased diagnostic testing in cattle.
- Separation of infected cattle during convalescence.
- Enforced cattle import restrictions to limit the movement of infected cattle across state borders.
- Stringent control of potentially infected food products (removal of milk or other infected products).

To minimize the potential for cow-to-human transmission, USDA recommends:

- Farm workers and other individuals in close contact with infected cattle or environments use adequate protective equipment and hygienic/sanitation measures.

Scenario 2: H5N1 virus stays predominantly in cattle but spreads widely within herds. There is low incidence of cow-to-human transmission but no human-to-human transmission.

In the second scenario, we considered the risk to human health if the H5N1 virus spreads widely within cattle herds but has minimal spread to other animal species. We determined the health risk to **farm workers** in this scenario to be **moderate**, the health risk to **other people in the vicinity of affected workers and animal populations** to be **low**, the risk to **healthcare workers** to be **low**, and the health risk to the **US general public** (and the consequent risk of a pandemic) to be **low**.

Our **confidence** in these risk scores is low, as the level of testing conducted in cattle and other farm animals is low. Based on available USDA data, there is considerable cow-to-cow transmission occurring. We do not know the level or types of exposure farm workers have to infected cattle, but it is likely sufficient to enable transmission. Furthermore, we do not have details on the transmission

HPAI A(H5) Scenario-Based Human Health Risk Assessment



routes to humans, making it difficult to determine the exact risk to workers who have contact with infected animals. Though we determined the risk of widespread human infection to be low given the low incidence of documented cow-to-human transmission, that could change. As the number of infected cattle increases, so does the risk to farm workers. To reduce risk, [USDA](#) recommends:

- Farm workers diligently use appropriate personal protective equipment (PPE; including masks, goggles, gloves, gowns, head covers, and boot covers) when working directly with or closely to cattle and poultry, other infected or potentially animals, and potentially infected environments.
- Increased diagnostic testing in cattle.
- Separation of infected cattle during convalescence.
- Enforced cattle import restrictions to limit the movement of infected cattle across state borders.
- Stringent control of potentially infected food products (removal of milk or other infected products).
- Information sharing between the agricultural and public health sectors to increase transparency and monitor for increases in animal-to-human transmission.

Scenario 3: H5N1 virus begins to infect swine or other animal species that could facilitate the mixing and spreading of influenza viruses. This increases the likelihood that the virus reassorts with other influenza viruses and adapts to humans. Some limited human-to-human transmission is reported, but only among close contacts of human cases. Healthcare workers have not reported infections.

In the third scenario, we considered the risk to human health if the H5N1 virus begins spreading widely in pigs or [other potential mixing vessel hosts](#), creating greater opportunity for reassortment with human influenza viruses. To date, naturally occurring H5N1 outbreaks have been reported in several of the potential mixing vessels, such as [minks, seals, cats, dogs, and many different kinds of birds](#). In this scenario, we expect some very limited human-to-human transmission would be reported among close contacts of cases. We determined the health risk to **farm workers** to be **moderate-high**, the health risk to **other people in the vicinity of affected workers and animal populations** to be **moderate**, the health risk to **healthcare workers** to be **low**, and the health risk to the **US general public** (and the consequent risk of a pandemic) to be **low**.

Our **confidence** in these risk scores is low, as the level of testing conducted among farm animals and people in contact with potentially infected animals is low. The jump from cattle to swine poses an increased risk for new mutations that could have a greater potential for sustained transmission within human populations, particularly for farm workers and local communities. To decrease risk to human health, we recommend:

HPAI A(H5) Scenario-Based Human Health Risk Assessment



- Farm workers diligently use personal protective equipment (PPE; including masks goggles, gloves, gowns, head covers, and boot covers) when working directly with or closely to cattle and poultry, other infected or potentially animals, and potentially infected environments.
- Increased diagnostic testing and genomic surveillance in cattle.
- Separation of infected cattle during convalescence.
- Enforced cattle import restrictions to limit the movement of infected cattle across state borders.
- Stringent control of potentially infected food products (removal of milk or other infected products).
- Information sharing between the agricultural and public health sectors to increase transparency and monitor for increases in animal-to-human or human-to-human transmission.
- Increased public health surveillance for H5N1 cases in local communities.
- Enhanced communication with the public about the situation and the measures being taken to address it.

Scenario 4: There are more reports of human infections due to contact with infected animals. Limited human-to-human transmission is reported among close contacts of infected individuals, including healthcare workers, but there is no efficient human-to-human transmission.

In the fourth scenario, we considered the risk to human health if the H5N1 virus begins spreading more readily among close human contacts, including healthcare workers. In this scenario, increased, but still limited, human-to-human transmission is reported among close contacts of infected individuals. Transmission between people is still not efficient. We determined the health risk to **farm workers** to be **high**, the health risk to **other people in the vicinity of affected workers and animal populations** to be **moderate-high**, the health risk to **healthcare workers** to be **moderate**, and the health risk to the **US general public** (and the consequent risk of a pandemic) to be **low-moderate**.

Our **confidence** in these risks scores is low, as the level of testing conducted among farm animals and humans is generally low. Increased incidence of human-to-human transmission may indicate increased transmission efficiency, but it may also be due to greater prevalence of the virus in communities. There is still significant uncertainty about whether the virus will spread efficiently among people. To reduce risk to human health, we recommend actions including but not limited to:

- Implementation of and support for recommended isolation of human cases and quarantine of close contacts of cases through escalated case finding and contact tracing, antiviral (eg, Tamiflu) prophylaxis for those exposed, compensation for individuals who are isolated/quarantined and cannot report to work, and social support to provide for essential needs of those in isolation/quarantine.

HPAI A(H5) Scenario-Based Human Health Risk Assessment



- Increased focus on sentinel surveillance, wastewater surveillance, and education of clinicians to consider H5N1 as a possible diagnosis for people who present with new respiratory illness.
- Continued development and widespread implementation of antigen and molecular testing in both hospital and outpatient healthcare settings.
- Policy preparation for the possibility of a pandemic, including congressional deliberations about emergency funding and emergency planning by healthcare institutions, workplaces, and federal, state, territorial, local, and tribal public health agencies.
- Increased investment and urgent development, testing, and production of vaccines and treatment options.
- Enhanced communication with the public about the situation and the measures being taken to address it, as well as efforts to mitigate the spread of rumors and disinformation.

Scenario 5: There are reports of efficient human-to-human transmission in addition to animal-to-human transmission. The likelihood of human infections is high because the virus now transmits efficiently and will be very difficult to contain.

In the fifth scenario, we considered the risk to human health if H5N1 continues to spread from animals to humans and develops the ability to transmit efficiently between humans. We determined the health risk to **farm workers** to be **high**, the health risk to **other people in the vicinity of affected workers and animal populations** to be **high**, the health risk to **healthcare workers** to be **high**, and the health risk to the **US general public** (and the consequent risk of a pandemic) to be **high**.

Our **confidence** in these risks scores is high. The level of H5N1 testing in humans is minimal, and we have limited information or evidence available to determine the exact human-to-human transmission pathways. Preventative medical countermeasures to address human-to-human transmission, such as vaccines, are available, but their effectiveness is unclear.

Nonpharmaceutical interventions are not currently in use, and there are no human public health response operations in place. The risk is therefore high for farm workers and the local and global communities. To decrease the risk of human-to-human transmission, we recommend actions including but not limited to:

- Increased diagnostic and surveillance testing in humans, including increased genomic surveillance.
- Implementation of and support for recommended isolation of cases and quarantine of close contacts of cases through escalated case finding and contact tracing, antiviral (eg, Tamiflu) prophylaxis for those exposed, compensation for individuals who are isolated/quarantined and cannot report to work, and social support to provide for essential needs of those in isolation/quarantine.

HPAI A(H5) Scenario-Based Human Health Risk Assessment



- Increasing sentinel surveillance, wastewater surveillance, and education of clinicians about how to recognize and treat H5N1 infection.
- Widespread implementation of antigen and molecular testing in both hospital and outpatient healthcare settings.
- Congressional approval of supplemental appropriations to fund public health response activities.
- Urgent development, testing, and production of vaccines and additional treatment options.
- Adoption and implementation of medical countermeasures and nonpharmaceutical interventions.
- Increased risk communication to the public to provide regular epidemiologic updates, discuss recommended and potential interventions, recommend measures that individuals and organizations can take to protect personal and public health, and address rumors and disinformation.
- Monitoring and mitigation of transmission at mass gathering events.

HPAI A(H5) Scenario-Based Human Health Risk Assessment



References

- Burrough ER, Magstadt DR, Petersen B, et al. Highly Pathogenic Avian Influenza A(H5N1) Clade 2.3.4.4b Virus Infection in Domestic Dairy Cattle and Cats, United States, 2024. *Emerg Infect Dis.* 2024;30(7):1335-1343. doi:10.3201/eid3007.240508
- Caserta LC, Frye EA, Butt SL, et al. Spillover of highly pathogenic avian influenza H5N1 virus to dairy cattle. *Nature.* July 25, 2025 (early release). doi:10.1038/s41586-024-07849-4
- Eisfeld AJ, Biswas A, Guan L, et al. Pathogenicity and transmissibility of bovine H5N1 influenza virus. *Nature.* July 8, 2024 (early release). doi:10.1038/s41586-024-07766-6
- Kristensen C, Jensen HE, Trebbien R, Webby RJ, Larsen LE. The avian and human influenza A virus receptors sialic acid (SA)- α 2,3 and SA- α 2,6 are widely expressed in the bovine mammary gland. *bioRxiv.* May 3, 2024 (early access). doi:10.1101/2024.05.03.592326
- Missouri Department of Health and Senior Services. Human H5 bird flu case confirmed in Missouri. Published September 6, 2024. Accessed October 3, 2024. <https://health.mo.gov/news/newsitem/uuid/0ca9d648-cb9b-4eca-99a8-f9a1c96a01bd/human-h5-bird-flu-case-confirmed-in-missouri>
- Schnirring, Lisa. CIDRAP. No Clear Exposure Source in Missouri H5 Avian Flu Case. September 24, 2024. Accessed October 3, 2024. <https://www.cidrap.umn.edu/avian-influenza-bird-flu/no-clear-exposure-source-missouri-h5-avian-flu-case>
- Science. ‘Incredibly concerning’: Bird flu outbreak at Spanish mink farm triggers pandemic fears. Published January 24, 2023. Accessed October 3, 2024. <https://www.science.org/content/article/incredibly-concerning-bird-flu-outbreak-spanish-mink-farm-triggers-pandemic-fears>
- Shittu I, Silva D, Oguzie J., et al. A one health investigation into H5N1 avian influenza virus epizootics on two dairy farms. July 31, 2024 (early release).doi:10.1101/2024.07.27.24310982
- University of California Davis. H5N1 is increasingly adapting to mammals: Elephant seal outbreak marks first transnational spread of highly pathogenic avian influenza in mammals. Published June 05, 2024. Accessed October 3, 2024. <https://www.ucdavis.edu/health/news/h5n1-increasingly-adapting-mammals#:~:text=They%20recorded%20unprecedented%20mass%20mortality,died%20at%20the%20same%20time>
- US Department of Agriculture Animal and Plant Health Inspection Service. Highly Pathogenic Avian Influenza (HPAI) Detections in Livestock. Updated August 30, 2024. Accessed October 3, 2024. <https://www.aphis.usda.gov/livestock-poultry-disease/avian/avian-influenza/hpai-detections/hpai-confirmed-cases-livestock>
- US Department of Agriculture Agricultural Research Service. Cow “Challenge” Study Should Help Turn Tables on H5N1 in Dairy Herds. Published July 17, 2024. Accessed October 3, 2024. <https://www.ars.usda.gov/news-events/news/research-news/2024/cow-challenge-study-should-help-turn-tables-on-h5n1-in-dairy-herds/>
- US Centers for Disease Control. CDC A(H5N1) Bird Flu response Update. Published July 24, 2024. Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/spotlights/h5n1-response-07192024.html#:~:text=CDC%20Recommendations&text=People%20should%20also%20avoid%20exposures,should%20not%20drink%20raw%20milk.>
- US Centers for Disease Control and Prevention. H5 Bird Flu: Current Situation. Updated

HPAI A(H5) Scenario-Based Human Health Risk Assessment



September 3, 2024. Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/situation-summary/index.html>

- US Centers for Disease Control and Prevention. CDC A(H5N1) Bird Flu Response Update. Published July 26, 2024 Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/spotlights/h5n1-response-07262024.html>
- US Centers for Disease Control and Prevention. CDC A(H5N1) Bird Flu response Update. Published July 19, 2024. Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/spotlights/h5n1-response-07192024.html>
- US Centers for Disease Control and Prevention. CDC A(H5N1) Bird Flu response Update. Published August 2, 2024. Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/spotlights/h5n1-response-08022024.html>
- US Centers for Disease Control and Prevention. CDC A(H5N1) Bird Flu response Update. Published September 13, 2024. Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/spotlights/h5n1-response-09132024.html>
- US Centers for Disease Control and Prevention. CDC A(H5N1) Bird Flu response Update. Published September 20, 2024. Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/spotlights/h5n1-response-09202024.html>
- US Centers for Disease Control and Prevention. CDC A(H5N1) Bird Flu response Update. Published September 27, 2024. Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/spotlights/h5n1-response-09272024.html>
- US Centers for Disease Control and Prevention. CDC Confirms Second Human H5 Bird Flu Case in Michigan; Third Case Tied to Dairy Outbreak. Published May 30, 2024. Accessed October 3, 2024. <https://www.cdc.gov/media/releases/2024/p0530-h5-human-case-michigan.html>
- US Centers for Disease Control and Prevention. CDC Confirms Human H5 Bird Flu Case in Missouri. Published September 6, 2024. Accessed October 3, 2024. <https://www.cdc.gov/media/releases/2024/s0906-birdflu-case-missouri.html>
- US Centers for Disease Control and Prevention. Update: Human Infection with Highly Pathogenic Avian Influenza A(H5N1) Virus in Texas. Published April 5, 2024 Accessed October 3, 2024. <https://www.cdc.gov/ncird/whats-new/human-infection-H5N1-bird-flu.html>
- US Centers for Disease Control and Prevention. 2024-2025 Influenza Season: Surveillance for Novel Influenza A and Seasonal Influenza Virus. Published September 20, 2024 Accessed October 3, 2024. <https://www.cdc.gov/bird-flu/php/monitoring-bird-flu/surveillance-novel-seasonal.html>
- Wolfe MK, Duong D, Shelden B, et al. Detection of hemagglutinin H5 influenza A virus sequence in municipal wastewater solids at wastewater treatment plants with increases in influenza A in spring, 2024. *Environ Sci Technol Lett*. 2024;11(6):526-532. doi:10.1101/2024.04.26.24306409
- UN Food and Agriculture Organization. Ongoing avian influenza outbreaks in animals pose risk to humans. Published July 12, 2023. Accessed October 3, 2024. <https://www.fao.org/animal-health/news-events/news/detail/ongoing-avian-influenza-outbreaks-in-animals-pose-risk-to-humans/en>